

UNDERSTANDING NATURAL VARIATIONS OF DISSOLVED METHANE IN AREAS OF ACCELERATING MARCELLUS SHALE GAS DEVELOPMENT

S. Sharma , M. Mulder, A. Sack , L. Bowman & T. Carr



K. Schroeder & R. Hammack




J. White & D. Chambers



DISSOLVED METHANE OR “STRAY GAS”

- ▣ Colorless & Odorless
- ▣ Flammable
- ▣ Explosive range: 5-15%
- ▣ Solubility in water : 26-32 mg/l





Problems:

Fires; Explosions; Asphyxiation; Groundwater Contamination

STUDY AREA

West Virginia Groundwater Well Site Locations:
within Marcellus 50 foot Isopach



Michon L. Mulder
West Virginia University




SAMPLED GW AQUIFERS

Dunkard Gp.
Monongahela Gp.
Conemaugh Gp.
Allegheny Fm.
Pottsville Gp.
Greenbrier Gp.
Pocono Gp.
Hampshire Fm.
Chemung Gp.



Sources of Stray Methane

- ▣ Landfills, Swamps and Marshes
- ▣ Microbial gas in shallow aquifers
- ▣ Abandoned & operating coal Mines
- ▣ Gas storage fields
- ▣ Gas pipelines
- ▣ Abandoned & operating gas wells



Pathways of Methane Formation

BIOGENIC

Bacterial gas

THERMOGENIC

Coal bed & Natural gas

ABIOGENIC

Crustal & mantle gas

ACETATE FERMENTATION




Near surface environment- Landfill, Marsh etc.

CO₂ REDUCTION




Drift gas from deeper formations



Using gas composition to identify methane origin



(Hunt, 1996)



Using stable isotopes to identify methane origin



$$\delta^{13}\text{C} (\text{‰}) = \left(\frac{{}^{13}\text{C}/{}^{12}\text{C}_{\text{sample}}}{{}^{13}\text{C}/{}^{12}\text{C}_{\text{standard}}} - 1 \right) \cdot 1000$$

$$\delta\text{D} (\text{‰}) = \left(\frac{{}^2\text{H}/{}^1\text{H}_{\text{sample}}}{{}^2\text{H}/{}^1\text{H}_{\text{standard}}} - 1 \right) \cdot 1000$$

Understanding Sources of Methane




GW methane data :




- MS thesis of M. Mulder

Methane data :


- Marcellus & Shallow Devonian Sands (samples collected)
- Ordovician & Silurian gases (published literature)





Other evidences of methane origin: $\delta^{13}\text{C}_{\text{DIC}}$



Relation to Mining and Abandoned Oil and Gas Activity



Relation to Topographic Curvature



Conclusions

Stable isotopic signatures and gas geochemistry can be used to

- ▣ Identify possible sources of methane in groundwater aquifers and surface waters
- ▣ Identify changes in hydrological connections

Essential criterion to be met:

- ▣ Good understanding of baseline and temporal variations in gas concentrations and isotopic signatures
- ▣ Well established dissolved gas sampling protocols

Ongoing Work...

- ▣ End-member characterization of natural gas in formations overlying Marcellus, active/in-active coal mines, microbial gas in shallow subsurface in southern Pennsylvania and north central West Virginia
- ▣ Assess effect of sampling methodologies on isotopic and molecular compositions of dissolved gases sampled using different techniques
- ▣ Test applicability of stable isotopes and gas composition to identify changes in hydrologic connections related to hydraulic fracturing.

ACKNOWLEDGMENTS

- URS -DOE National Technology Laboratory
- US Geological Survey
- West Virginia Water Research Institute
- USGS West Virginia Water Science Center